

Intended use

Eosin (Aqu.) 2% is used as monochrome for bacterial staining and as a counter stain for hematoxylin.

Summary

Eosin is most often used as a counterstain to hematoxylin in H&E (hematoxylin and eosin) staining. Eosin is a name of several fluorescent acidic compounds which bind to and from salts with basic, or eosinophilic, compounds like proteins containing amino acid residues such as arginine and lysine and stains them dark red or pink because of the actions of bromine on fluorescein. In addition to staining proteins in the cytoplasm, it can be used to stain collagen and muscle fibers for examination under the microscope. Structures that stain readily with eosin are termed eosinophilic.

Principle

Hematoxylin and eosin are the principal stains used for the demonstration of nucleus and the cytoplasmic inclusions. Alum acts as a mordant and hematoxylin containing alum stains nucleus light blue which turns red in the presence of acid. The cell differentiation is achieved by treating the tissue with acid solution. Counter staining is performed by using eosin solution which imparts pink color to the cytoplasm.

Hematoxylin, a common nuclear stain, is isolated from an extract of logwood (*Haematoxylon campechianum*). Before hematoxylin can be used as a nuclear stain, it must be oxidized to hematein and combined with a metallic ion (mordant). Most successful mordants have been salts of aluminum or iron. Generally, hematoxylin are classified as progressive or regressive based on dye concentration. Progressive stains (e.g., Mayer's hematoxylin) have a lower concentration of dye and selectively stain nuclear chromatin. The desired intensity is a function of time. Regressive stains (e.g., Harris hematoxylin) color all nuclear and cytoplasmic structures intensely. To arrive at correct chromatic response, excess dye must be removed by treatment with dilute acid (differentiation).

Eosin is tetra bromofluorescein (a substituted xanthene), a red acidic dye and fluorochrome. The dye is very soluble in ethyl alcohol and also used for the staining of cytoplasm. Eosin Y is the most commonly used counterstain for hematoxylin.

Reagents / Contents

Eosin (AQU.) 2%

Eosin-Y 2.0 g

Distilled water 100 mL

Appearance: Dark reddish colour solution.

Storage and Stability

Store at 15°C-25°C away from bright light. Use before expiry date on label.

Materials required but not provided

Tissue section specimen on clean grease-free glass slide, staining rack, blotting paper, immersion oil, and microscope. Reagents and solutions required but not provided in the Kit such as xylene, a series of descending and of ascending grades of alcohol, 1% acid alcohol solution, Scott's Tap Water Buffer, Hematoxylin Harris and DPX mountant.

Type of Specimen

Histochemical tissues sections obtained from biopsy specimens.

Procedure

1. Sections are deparaffinized (removal of wax) by placing in xylene for 10 - 15 minutes.
2. Rehydrate section by passing in a series of descending grades of alcohol, finally to water.
3. Place in Hematoxylin Harris solution for 8-10 minutes.
4. Rinse in water.
5. Differentiate the slide in a solution 1% acid alcohol for 10 seconds.
6. Rinse in tap water.
7. Blueing (bringing the required blue color to section) is done by putting the section in a solution containing Sodium bicarbonate, MgSO₄ and saturated solution of Lithium carbonate (Scott's Tap Water Buffer) for 2-10 minutes.
8. Counter stain with aqueous Eosin (Aqu.) 2% for 1-3 minutes.
9. Rinse in tap water.
10. Section are dehydrated which is done by a series of ascending grades of alcohol and finally clearing in Xylene.
11. Dry the section by pressing on the filter paper.
12. Mount in DPX mountant and observe under microscope, 40X and 100X under oil immersion lens.

Interpretation of Results

The nuclei of cells are stained blue or dark purple along with a few other tissues, such as keratohyalin granules and calcified material with Hematoxylin. The cytoplasm and some other structures including extracellular matrix such as collagen stains in up to five shades of pink with Eosin. Most of the cytoplasm is eosinophilic and is rendered pink. Red blood cells are stained intensely red. The background of the tissue remains colourless.

Warranty

Eosin (Aqu.) 2% staining solution is for "In Vitro Diagnostic Use" only. This product is designed to perform as described on the label and pack insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

Data on file: UltraCare Diagnostics .

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.